

СОДЕРЖАНИЕ

Обзоры

<i>Martell J. M., Mitrea D., Mitrea I., Mitrea M.</i> Fatou-type theorems and boundary value problems for elliptic systems in the upper half-space . . .	3
<i>Tkachev V. G.</i> Spectral properties of nonassociative algebras and breaking regularity for nonlinear elliptic type PDEs	51

Статьи

<i>Akman M., Lewis J., Vogel A.</i> Note on an eigenvalue problem for an ODE originating from a homogeneous p -harmonic function	75
<i>Алхутюв Ю. А., Сурнячѐв М. Д.</i> Поведение в граничной точке решений задачи Дирихле для $p(x)$ -лапласиана	88
<i>Arkhipova A. A.</i> Weak global solvability of the two-phase problem for a class of parabolic systems with strong nonlinearity in the gradient. The case of two spatial variables	118
<i>Асташиова И. В.</i> О петичности асимптотически степенных решений уравнения типа Эмдена–Фаулера высокого порядка	152
<i>Brennan J. E.</i> Bounded point derivations on certain function spaces	174
<i>Gol'dshstein V., Pchelintsev V., Ukhlov A.</i> On conformal spectral gap estimates of the Dirichlet-Laplacian	189
<i>Davey B., Kenig C., Wang J.-N.</i> On Landis' conjecture in the plane when the potential has an exponentially decaying negative part	204
<i>Lanzara F., Schmidt G.</i> Approximate approximations: recent developments in the computation of high dimensional potentials	227
<i>Miyazishi Y., Rozenblum G.</i> Eigenvalues of the Neumann–Poincare operator in dimension 3: Weyl's law and geometry	248
<i>Seregin G., Wang W.</i> Sufficient conditions on Liouville type theorems for the 3D steady Navier–Stokes equations	269